

**Original citation:**

Hong, Yiguo and Jackson, Stephen D.. (2015) Floral induction and flower formation : the role and potential applications of miRNAs. Plant Biotechnology Journal . ISSN 1467-7644

**Permanent WRAP url:**

<http://wrap.warwick.ac.uk/66841>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.


**Publisher's statement:**

This is the accepted version of the following article: Hong, Yiguo and Jackson, Stephen D.. (2015) Floral induction and flower formation : the role and potential applications of miRNAs. Plant Biotechnology Journal . ISSN 1467-7644, which has been published in final form at: <http://onlinelibrary.wiley.com/doi/10.1111/pbi.12340/abstract>

**A note on versions:**

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP url' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: [publications@warwick.ac.uk](mailto:publications@warwick.ac.uk)



<http://wrap.warwick.ac.uk>

# **Floral induction and flower formation -the role and potential applications of miRNAs.**

Yiguo Hong<sup>1</sup> and Stephen Jackson\*

School of Life Sciences, Warwick University, Coventry, CV4 7AL, UK

<sup>1</sup>Research Centre for Plant RNA Signalling, College of Life and Environmental Sciences, Hangzhou Normal University, Hangzhou 310036, China.

\* Corresponding author

Tel: +44 (0)24 76522571

e-mail: [Stephen.jackson@warwick.ac.uk](mailto:Stephen.jackson@warwick.ac.uk)

Y. Hong's e-mail: [yiguo.hong@warwick.ac.uk](mailto:yiguo.hong@warwick.ac.uk)

Running title: Role of miRNAs in flowering

Keywords: Flowering time, Flower development, Juvenility, Floral induction, Floral patterning, male sterility, miRNA

Word count: 7,061 (without bibliography)

The authors declare that they have no conflict of interest

## **Abstract**

The multiple regulatory pathways controlling flowering and flower development are varied and complex, and they require tight control of gene expression and protein levels.

MicroRNAs (miRNAs) act at both the transcriptional and post-transcriptional level to regulate key genes involved in flowering-related processes such as the juvenile-adult transition, the induction of floral competence, and flower development. Many different miRNA families are involved in these processes and their roles are summarized in this review, along with potential biotechnological applications for miRNAs in controlling processes related to flowering and flower development.

## **Introduction**

There are a number of different developmental phases through which a plant progresses during its life-cycle, these are differentiated by distinct morphological traits and/or the development of new organs (Huijser and Schmid, 2011; Jin et al., 2013). Flowering is an essential part of the reproductive process in angiosperms, and the flowering process involves two developmental phase transitions: the first of these is the transition from the immature juvenile vegetative state (where the plant is unable to flower) to the mature adult vegetative state (where the plant is capable of flowering) (Poethig 2003; Sgamma et al., 2014). The second developmental transition occurs at floral induction when the plant

switches from vegetative growth to reproductive growth and starts to produce flowers. The timing of this transition is tightly controlled by a complex gene regulatory network, this is because the time when a plant flowers affects its reproductive success, eg. flowering usually occurs at the optimum time for pollination or seed dispersal. For out-crossing species this is of particular importance as flowering time needs to be co-ordinated with other plants of the same species in their vicinity, and for insect pollinated plants with the activity of their pollinators (Huijser and Schmid, 2011; Srikanth and Schmid, 2011). To make this possible the plant detects environmental cues such as light quality and duration, and temperature, and integrates this information with that from endogenous physiological processes such as its circadian clock, phytohormone and carbohydrate levels, and vernalisation state in order to regulate its flowering time to the appropriate time of the year, and even time of the day (Amasino 2010; Fornara et al., 2010; Jackson 2009; Kim et al., 2012; Srikanth and Schmid, 2011; Matsoukas et al., 2012).

As the timing of flowering and floral organogenesis both significantly affect plant fitness and crop yield, a detailed understanding of the regulatory mechanisms governing flowering time is essential for continued improvements in agricultural practice (Huijser and Schmid, 2011; Srikanth and Schmid, 2011). A lot of research has therefore been conducted on the molecular mechanisms controlling the flowering process, and numerous genes that contribute to the different environmental and endogenous regulatory pathways have been identified. In *Arabidopsis*, it is now well known that the main pathways controlling flowering in response to environmental signals are the photoperiod, ambient temperature and vernalization pathways, which respond to day length, surrounding temperature and

prolonged cold exposure, respectively (Fornara et al. 2010; Jackson 2009). Other endogenous factors such as phytohormones and carbohydrate status also regulate flowering through the autonomous, gibberellic acid (GA), nutrient-responsive and aging pathways (Kim et al., 2012; Srikanth and Schmid, 2011; Matsoukas et al., 2012; Yamaguchi and Abe, 2012). All these different regulatory pathways converge on a set of floral pathway integrator genes, namely *FLOWERING LOCUS T* (*FT*) and its paralogue *TWIN SISTER OF FT* (*TSF*), as well as *SUPPRESSOR OF CONSTANS 1* (*SOC1*) and *AGAMOUS-LIKE 24* (*AGL24*). These act to control the expression of a small set of meristem identity genes at the shoot apical and lateral meristems including *LEAFY* (*LFY*), *APETALA 1* (*API*) and *FRUITFUL* (*FUL*). Once the expression of these genes reaches a certain level they induce the expression of floral organ identity genes and flowers are produced (Amasino 2010; Huijser and Schmid, 2011; Pose et al., 2012; Srikanth and Schmid, 2011).

Over the past decade there have been numerous reports demonstrating that a number of different microRNA (miRNA) families play important roles in the regulation of flowering time and floral development, which they do by targeting and down-regulating transcription factors involved in these processes (Chuck et al., 2009; Jones-Rhoades et al., 2006; Luo et al., 2013; Spanudakis and Jackson 2014). Much of this evidence has been obtained from research on the model plant *Arabidopsis*, although a high level of conservation of miRNAs and their regulatory pathways/target genes has been demonstrated from results obtained from tomato, potato, maize, rice, and other species (Willmann and Poethig, 2007; Sunkar and Jagadeeswaran, 2008; Luo et al., 2013). There are at least eleven different miRNA families that have been shown to affect different stages in the induction of flowering and

flower development in plants, these are *miR156*, *miR159*, *miR160*, *miR164*, *miR166/165*, *miR167*, *miR169*, *miR172*, *miR319*, *miR390* and *miR399* (see Fig.1).

## **Regulation of the time of flowering**

The main miRNA families involved in regulating flowering time are the *miR156*, *miR172* and *miR390* families which are involved in the juvenile-to-adult vegetative phase transition, and the *miR159*, *miR169* and *miR399* families which along with *miR156* and *miR172* are involved in the vegetative-to-reproductive phase transition (Huijser and Schmid, 2011; Jin et al., 2013; Jones-Rhoades et al., 2006; Kim et al., 2011; Rubio-Somoza and Weigel, 2011; Wang 2014; Zhu and Helliwell, 2011).

### The role of *miR156*

In Arabidopsis, the *miR156* family is encoded by the loci *MIR156a-j* (Yamaguchi and Abe, 2012). The *miR156* family acts together with the *miR172* family within the ageing pathway to regulate the time when a plant becomes mature and reproductively competent (Huijser and Schmid, 2011; Yamaguchi and Abe, 2012; Wang et al., 2009; Wu et al., 2009). *MIR156* regulates the levels of *miR172* such that they have temporally opposite expression patterns, *miR156* is highly expressed in the embryo and early seedling stage and declines with increasing plant age whereas *miR172* accumulates over time in the leaves and floral buds (Fahlgren et al., 2006; Wu et al., 2009; Zhu and Helliwell, 2010). This forms the basis for the regulation of the juvenile to adult phase change and, in addition to flowering

time, other morphological traits representative of the juvenile and adult vegetative phases such as leaf morphology and trichome distribution are also affected by the activity of these miRNAs (Huijser and Schmid, 2011).

Transgenic plants constitutively overexpressing *miR156* exhibited delayed-flowering and a prolonged juvenile phase, as shown by the increased production of juvenile leaves and lack of abaxial trichomes (Huijser and Schmid, 2011; Wu and Poethig, 2006). This effect was more pronounced at 16°C than at 23°C leading to the suggestion that ambient temperature may influence the effect of *miR156* over-expression (Kim et al., 2012). An evolutionary conserved role of *miR156* in the control of flowering is supported by the fact that delayed flowering is also observed in rice, tomato and maize lines over-expressing *miR156* (Xie et al., 2006; Zhang et al., 2011; Chuck et al., 2007). On the other hand, expression of a target mimic of *miR156* (*MIM156*), which sequesters the available *miR156* and down-regulates *miR156* activity, resulted in early flowering after producing very few leaves that all had adult features (Franco-Zorrilla et al., 2007, Todesco et al., 2010). Thus, high levels of *miR156* early in plant development suppress flowering and are necessary for the expression of the juvenile phase. (Huijser and Schmid, 2011).

The targets of the *miR156* family are the SQUAMOSA PROMOTER BINDING-LIKE (SPL) transcription factors, *miR156* down-regulating the expression levels of 11 out of the 17 *SPL* genes in Arabidopsis (Franco-Zorrilla et al., 2007; Huijser and Schmid, 2011; Yamaguchi and Abe, 2012). The decline in *miR156* levels over time with increasing age is therefore accompanied by a corresponding increase in the expression level of *SPL* transcription factors which promote flowering through the induction of *FT*, *LFY* and *MADS*-box gene expression (Wu and Poethig, 2006; Yamaguchi and Abe, 2012). There is

functional redundancy within the *SPL* gene family, and the loss of one *SPL* protein often has no effect (Yamaguchi and Abe, 2012). The *SPL* genes have therefore been grouped together into four clades, two of which (clade VI and clade VIII) have a significant influence on flowering (Guo et al., 2008; Huijser and Schmid, 2011; Wu et al., 2009). Clade VIII comprises *SPL9* and *SPL15* which act redundantly, with *spl9 spl15* double loss-of-function mutants exhibiting a similar phenotype to plants overexpressing *miR156* (Guo et al., 2008; Schwarz et al., 2008). On the other hand, plants expressing *miR156*-resistant forms of *SPL9* or *SPL15* (*rSPL9* or *rSPL15*), that are therefore not down-regulated by *miR156*, flower very early due to the induction of *miR172* expression (Wu et al., 2009; Zhu and Helliwell, 2010). *SPL9* has been shown to directly bind to, and activate the transcription of, *miR172*, and it has also been shown to be a direct regulator of *SOC1*, *AGL24*, *FUL* and *AP1* (Wang et al., 2009).

Clade VI consists of the *SPL3*, *SPL4* and *SPL5* genes (Huijser and Schmid, 2011). The expression of *miR156*-resistant *rSPL3*, *rSPL4*, or *rSPL5* genes, which lack the *miR156* binding site in the 3' UTR, resulted in increased levels of *SPL* proteins and premature appearance of adult leaf traits and early flowering (Guo et al. 2008; Huijser and Schmid, 2011; Wu et al., 2009). *SPL3* has been shown to bind the promoter and intragenic elements of the floral meristem identity genes *LFY*, *FUL* and *AP1*, and so appears to target the same floral meristem identity genes as *SPL9* (Yamaguchi et al., 2009).

The main role of *miR156* is thus to prevent flowering until plants have reached a certain developmental stage, ie. until they have completed their juvenile phase. This is evidenced by the fact that *miR156* has been shown to prevent precocious flowering (Wang et al., 2009). It inhibits flowering by i). repressing the expression of *SPL* genes which are



direct activators of floral integrator and floral meristem identity genes, and ii). repressing the induction of *miR172* and in doing so maintaining high levels of expression of AP2-like floral repressors which repress flowering (see below).

In this way the plant is not ‘competent’ to respond to inductive environmental signals and cannot be induced to flower whilst high levels of *miR156* are present (ie. during the juvenile phase). As the plant grows older and the levels of *miR156* decline then this repression of flowering is gradually reduced and the plant becomes capable of responding to environmentally inductive signals (ie. the plant has become mature).

What causes the reduction in *miR156* levels with age is not well understood. The findings that metabolically active sugars such as glucose and sucrose, or the levels of trehalose-6-phosphate (T6P) which serves as a signal for carbohydrate availability in the plant, selectively regulate the expression of *miR156* genes however, suggests that nutritional status may serve as a signal for the age or developmental stage of the plant (Wahl et al., 2013; Yang et al., 2013; Yu et al., 2013). Sugar accumulation reduces *miR156* levels and conversely sugar deprivation increases *miR156* levels with a corresponding increase or decrease in SPL levels respectively. At least some of the effects of sugar levels on *miR156* levels are mediated by the glucose sensing enzyme HEXOKINASE 1 (HXK1) which up-regulates *miR156* levels in low sugar conditions (Yang et al., 2013). Another link between carbohydrate levels and *miR156* expression has been shown to be mediated by T6P (Wahl et al., 2013). In transgenic lines with reduced levels of T6P, due to knock-down of *TREHALOSE-6-PHOSPHATE SYNTHASE 1 (TPS1)* expression, flowering was delayed, and in younger plants *miR156* levels were higher and *SPL3*, *SPL4* and *SPL5* expression

levels were correspondingly reduced. The levels of *miR156* still decreased with age in these plants but as they were initially at a higher level they took longer to decline (Wahl et al., 2013).

*MiR156* levels have also shown to be affected by cold temperatures in the perennial plant *Arabis alpina*, a relative of *Arabidopsis* that requires vernalisation to be induced to flower. Exposing *A.alpina* seedlings to prolonged periods of cold delayed the age-dependent decline of *miR156* levels thus increasing the age at which the plants could respond to vernalization, the cold temperature was shown to regulate transcription of the *MIR156* gene (Bergonzi et al., 2013).

#### The role of *miR172*

The *miR172* family, which are coded by the *MIR172a-e* loci, have the opposite effect on flowering time to *miR156*; overexpression of *miR172* (*35S::miR172b*) results in an extremely early flowering phenotype in both inductive long day (LD) and non-inductive short day (SD) conditions (Aukerman and Sakai, 2003; Wu et al., 2009; Zhu and Helliwell, 2010). *MiR172* acts downstream of *miR156* and the *SPL9*, *10*, *11* and *15* genes, the latter acting redundantly to regulate *miR172* levels (Wu et al., 2009). There are multiple copies of *SPL* binding elements in the promoter of the *MIR172b* gene, and chromatin immunoprecipitation showed that at least one is strongly bound by *SPL9*, furthermore plants over-expressing *SPL9* were found to have increased levels of *miR172*. The increase in *miR172* levels with age (a temporally opposite expression pattern to *miR156*) is thus a direct consequence of reducing *miR156* levels and increased expression of *SPL* genes (Wu et al., 2009).

In *Arabidopsis* *miR172* targets the expression of a family of AP2-like transcription factors including APETELA 2 (AP2), TARGET OF EAT 1 (TOE1), TOE2, TOE3, SCHLAFMÜTZE (SMZ) and SCHNARCHZAPFEN (SNZ) (Aukerman and Sakai, 2003; Chen, 2004; Yamaguchi and Abe, 2012). These AP2-like proteins are floral repressors and delay flowering by inhibiting expression of *FT* and floral meristem identity genes *AP1*, *LFY*, *FUL* and *SOC1* (Mathieu et al., 2009; Yant et al., 2010; Zhu and Helliwell, 2010). Overexpression of AP2-like genes such as *SMZ* and *SNZ* results in a late flowering phenotype (Mathieu et al., 2009; Yamaguchi and Abe, 2012). AP2-type protein levels are high in the early seedling and juvenile stage and decline as *miR172* levels rise with increasing plant age thus relieving the repression of flowering as the plant matures (Jung et al., 2007; Zhu and Helliwell, 2010). AP2 acts in a feedback loop which up-regulates *miR156* levels and down-regulates *miR172* levels and is thought to fine-tune the flowering response (Yant et al., 2010; Huijser and Schmid, 2011). Furthermore AP2-type proteins regulate the expression of other AP2-type genes which adds another layer of feedback regulation (Zhou and Wang, 2013; Zhu and Helliwell, 2010).

In addition to the aging pathway *miR172* is also regulated by other flowering pathways such as the photoperiod, ambient temperature and GA-regulated flowering pathways (Jung et al., 2007; Lee et al., 2010; Yamaguchi and Abe, 2012). In the photoperiodic pathway, GI up-regulates *miR172* in a CO-independent manner (Yamaguchi and Abe, 2012; Zhou and Wang, 2013). GI affects the processing of *miR172* rather than its transcription because in the *gi* mutant levels of *miR172* are reduced despite increased levels

of the primary *MIR172* (*pri-MIR172*) transcript (Jung et al., 2007). In the ambient temperature pathway, the floral repressor SHORT VEGETATIVE PHASE (SVP), and the RNA-binding protein FCA both repress the expression of *miR172* (Kim et al., 2012; Zhu and Helliwell, 2010). Low ambient temperatures increase the levels of SVP and FCA therefore resulting in decreased *miR172* levels. Loss of *SVP* results in ambient temperature insensitive flowering (Kim et al., 2012; Yamaguchi and Abe, 2012). In the GA pathway DELLA proteins repress flowering in part through repressing SPLs and *miR172* levels, and thus the induction of *FT*, in leaves (Galvao et al., 2012; Yu et al., 2012). *MiR172* and its target genes are thus involved in several different flowering time pathways and this miRNA therefore represents a hub for the integration of these separate pathways. There is some evidence that *miR172* can move across graft unions and throughout the plant suggesting that it may play a role as a systemically mobile developmental signal (Martin et al., 2009; Kasai et al., 2010).

### The role of *miR390*

*MiR390* is involved in multiple developmental processes including leaf morphogenesis, lateral root development, and flowering time control. Its effects on flowering time are because it prolongs the juvenile phase and thus delays the acquisition of the competence to flower (Fahlgren et al., 2006; Rubio-Somoza and Weigel, 2011).

*MiR390* mediates its effects on flowering time not by directly targeting protein-coding mRNAs but by triggering the production of trans-acting siRNAs from the *TAS3* locus, which in turn target and repress mRNA levels of the transcription factors AUXIN RESPONSE FACTORS 3 (ARF3) and ARF4 (Endo et al., 2013; Garcia, 2008;

Montgomery et al., 2008; Rubio-Somoza and Weigel, 2011). ARF3 and ARF4 activity promotes the juvenile-to-adult vegetative phase transition and mutants defective in tasiRNA biogenesis have a shorter juvenile phase as a result of elevated levels of ARF3 and ARF4. Transgenic plants expressing tasiRNA-insensitive *ARF3* (*ARF3:ARF3mut*) also exhibited a shorter juvenile phase (Fahlgren et al., 2006; Garcia, 2008). *MiR390* thus delays flowering by repressing ARF3 and ARF4 activity and prolonging the juvenile phase (Fahlgren et al., 2006; Garcia, 2008; Rubio-Somoza and Weigel, 2011).

ARF3 and ARF4 may affect the juvenile-to-adult transition by affecting the expression of *miR156*-regulated *SPL* genes (Rubio-Somoza and Weigel, 2011), and as AP2 (a target of *miR172*) has been shown to directly bind to the *ARF3* promoter to repress its expression (Yant et al., 2010), there is therefore a link between *miR390* and the *miR156/miR172* feedback interaction in the regulation of juvenile phase length and the juvenile-to-adult transition (Rubio-Somoza and Weigel, 2011).

### The role of *miR159*

*MiR159* affects the time of flowering through its role in the GA regulation of floral induction, and is also part of a network involving two other miRNAs, *miR319* and *miR167*, that controls floral organ development (see Fig.3 and below). In *Arabidopsis* the *miR159* family is encoded by three loci (*MIR159a-c*), and are involved in the GA flowering pathway, which promotes flowering under non-inductive SD conditions (Terzi and Simpson, 2008; Yamaguchi and Abe, 2012). DELLA proteins, which repress the GA response, also repress the expression of *miR159* and its target genes, *MYB33*, *MYB65* and *MYB101*, which are homologues of the rice and barley GAMYB transcription factors

(Achard et al., 2004; Allen et al., 2007). GA treatment results in the degradation of DELLA proteins with a resulting increase in both *miR159* levels and levels of the GAMYB transcription factors which then are able to bind to GA-response elements in the *LFY* promoter to induce its transcription (Achard et al., 2004; Jin et al., 2013). As putative GA-response elements have also been identified in the *miR159* promoter it is possible that the GAMYB factors may also enhance expression of *miR159* in a feedback loop that will then down-regulate GAMYB expression to provide a level of homeostatic regulation of the GA response (Fig 3). This is supported by the fact that overexpression of *miR159* causes a decrease in *MYB33* and *LFY* transcript levels and a delay in the onset of flowering (Achard et al., 2004; Li et al., 2013).

### The role of *miR169*

The *miR169* family is the largest miRNA family. In Arabidopsis is made up of 14 members *miR169a-n* (Li et al., 2010), and they target the *NF-YA* transcription factor gene family which is involved in the transcriptional regulation of a large number of genes (Jones-Rhoades and Bartel, 2004; Zhao et al., 2009). *MiR169* is an example of a miRNA involved in both abiotic stress response and the control of flowering time, most members of the *miR169* families in Arabidopsis, maize, and soybean are up-regulated by abiotic stress. It has been shown that over-expression of *miR169d* in Arabidopsis results in early flowering, and conversely over-expression of a *miR169d*-resistant version of *NF-YA2*, which is a *miR169d* target gene, causes late flowering (Xu et al., 2013). The effect on flowering time was shown to be mediated through the regulation of *FLC* expression by NF-

YA2 which binds to the promoter and first intron of the *FLC* gene to induce its expression. The reduced levels of NF-YA2 in *miR169d* over-expressing plants results in reduced *FLC* expression and increased expression of *FT* and *LFY* and therefore early flowering. The opposite happens in plants over-expressing the *miR169d*-resistant NF-YA2 where *FLC* expression is increased and *FT* and *LFY* expression levels are reduced resulting in late flowering (Xu et al., 2013). These results suggest that *miR169* might play a key role in the promotion of early flowering by abiotic stresses.

### The role of *miR399*

In *Arabidopsis* the *miR399* family is encoded by the *MIR399a-f* loci. It is a key player in phosphate homeostasis, which it affects through its down-regulation of the expression of *PHOSPHATE 2 (PHO2)* (Kim et al., 2011; Kruszka et al., 2012). PHO2 is an E2 ubiquitin-conjugating enzyme that targets proteins involved in phosphate uptake in the roots (Kim et al., 2011; Liu and Vance, 2010). *MiR399* activity is up-regulated under phosphate starvation to increase phosphate uptake, and down-regulated under high phosphate conditions to avoid phosphate toxicity, its activity being tightly controlled to prevent excessive phosphate accumulation and tissue necrosis (Chiou et al., 2006; Liu and Vance, 2010; Kruszka et al., 2012).

A potential role for *miR399* as an ambient temperature-responsive flowering time regulator has been suggested (Kim et al., 2011). Plants flower earlier when grown at 23°C than 16°C, and *miR399* levels were shown to be higher in plants grown at 23°C than those grown at 16°C. Furthermore, both *miR399* overexpressors and *PHO2* loss-of-function mutants flowered earlier than wild-type when grown at 23°C whereas this difference was

not observed when they were grown at 16°C. The earlier flowering of these plants at 23°C could be explained by increased levels of *TSF* expression (Kim et al., 2011). *MiR399* is thus another example of a miRNA involved in both abiotic stress response and the control of flowering time.

### **Regulation of floral patterning and floral organ development**

Once the plant is induced to flower the floral meristem genes induce the expression of floral organ identity genes which direct the formation of floral organ primordia at the meristem. The formation of floral organs occurs in successive whorls which in *Arabidopsis* is a whorl of four sepal primordia, a whorl of four petal primordia, a whorl of six stamen primordia, and finally two carpel primordia. These distinct whorls require boundaries to be established that limit the expression of floral organ identity genes as per the ABC model (Causier et al. 2010). MiRNAs such as *miR164*, *miR169* and *miR172* are involved in setting up such boundaries. Other miRNAs are involved in the subsequent development of floral organs such as petals, sepals, anthers and carpels. These include *miR159*, *miR160*, *miR165*, *miR166*, *miR167*, and *miR319*, some of which act together in regulatory modules (Fig.1; Rubio-Somoza and Weigel, 2013).

#### The role of *miR164* in flower development

The *miR164* family is made up of three members *miR164a-c*. This family down-regulates the expression of members of the NAC-domain family of transcription



factors such as the Arabidopsis *CUP-SHAPED COTYLEDON1* (*CUC1*) and *CUC2* genes, and the palm *NO APICAL MERISTEM* genes *EgNAM1* and *PdNAM1* (Takada et al., 2001; Vroemen et al., 2003; Adam et al., 2011). These transcription factors are involved in defining morphogenetic boundaries in the establishment of the floral meristem. Reducing the activity of the *CUC1* and *CUC2* genes changes the sepal boundary and results in fused sepals and a reduced number of petals, and *miR164c* has been shown to control petal number by regulating the expression levels of *CUC1* and *CUC2* (Laufs et al., 2004; Baker et al., 2005). *MiR164* has also been shown to be involved in setting boundaries between the SAM and leaf primordia in tomato, where it has a complimentary pattern of expression in the SAM to its NAC transcription factor target *GOBLET* (*GOB*). Mis-expression of *miR164* or expression of a *miR164*-insensitive *GOB* gene results in defective leaf development (Berger et al., 2009). In addition to its role in setting the sepal-petal and leaflet boundaries, *miR164* has been shown to be involved in carpel development, with Arabidopsis mutants of *miR164* exhibiting defects in carpel closure (Baker et al., 2005; Sieber et al., 2007).

#### The role of *miR169* in flower development

In *Petunia* and *Antirrhinum* *miR169* family members target the *NF-YA* transcription factor gene family which are involved in the activation of the homeotic C-class genes in flower development (Jones-Rhoades and Bartel, 2004; Zhao et al., 2009). These *miR169* genes are called *BLIND* (*BL*) in *Petunia* and *FISTULATA* (*FIS*) in *Antirrhinum*, and they regulate *NF-YA* gene expression to restrict the activity of the C-

class genes to the inner two floral whorls during flower development. Loss-of-function *fis* and *bl* mutations result in conversion of petals to stamens in the second whorl, indicating that there is abnormal C-gene activity in the second whorl due to the lack of *miR169* (Cartolano et al., 2007). Whilst such a role for *miR169* in floral development has been shown in *Petunia* and *Antirrhinum*, this does not appear to be the case in *Arabidopsis*, however, where *miR169* does not regulate C-gene activity (Cartolano et al., 2007).

#### The role of *miR172* in flower development

*MiR172* plays a role in floral patterning because it determines the domain of expression of the floral organ identity gene *AP2*. *MiR172* acts at the post-transcriptional level to restrict *AP2* expression to the two outer whorls of the floral meristem which give rise to sepals and petals. The domains of expression of *miR172* in the inner two whorls and *AP2* expression in the outer two whorls are largely complimentary as shown by *in situ* hybridization (Chen, 2004; Zhao et al., 2007; Wollmann et al., 2010), however at some timepoints an overlap in expression domains has been observed suggesting that other factors in addition to *miR172* may be regulating the extent of *AP2* expression (Wollmann et al., 2010). There is also evidence of a positive feedback loop whereby *AP2* is involved in the repression of *miR172* expression in the outer whorls. Yant et al. (2010) identified *MIR172b* as a target of *AP2* using ChIP-seq, and Grigorova et al. (2011) showed that the transcriptional repressor LEUNIG (*LUG*) directly represses *miR172* expression in sepals and that this repression requires the presence of *AP2*. It is proposed that *AP2* is responsible

for recruiting the LUG transcriptional repressor complex to the *MIR172* genes to repress *miR172* levels in the outer floral whorls and thus maintain its own expression in those whorls (Grigorova et al., 2011).

*MIR172* has been shown to be involved in floral patterning in cereals such as rice and barley where it targets the AP2-like transcription factor genes *SUPERNUMARY BRACT (SNB)* and *Oryza sativa INDETERMINATE SPIKELET1 (OsIDS1)* in rice, and *CLEISTOGAMY1 (Cly1)* in barley, to regulate lodicule development (Lee and An, 2012; Nair et al., 2010; Zhu et al., 2009).

#### The roles of *miR159*, *miR167* and *miR319* in flower development

*MIR159* and *miR319* have overlapping roles in controlling floral development as the target genes they regulate, the MYB and TCP transcription factors respectively, interact with each other and both act to regulate levels of *miR167* (Fig.3). *MIR159*, *miR319* and *miR167* therefore act together forming a regulatory circuit. Despite the fact that *miR159* and *miR319* evolved from a common ancestor and have 17 identical nucleotides (Li et al., 2011), these miRNAs do not cross-regulate each other's target genes. *MIR159* cannot bind *TCP* transcripts, and whilst *miR319* is capable of binding *MYB* transcripts, it has a much more limited spatial and temporal expression pattern compared to *miR159*. Because of this, in addition to their overlapping functions *miR159* and *miR319* can also play distinct regulatory roles in plant development (Jones-Rhoades et al., 2006; Palatnik et al., 2007).

As mentioned previously *miR159* targets *GAMYB*-related genes which, in addition to regulating *LFY* transcript levels and thus impacting on flowering time, are also involved

in anther development (Achard et al., 2004). *MiR159* restricts the expression of *GAMYB* genes such as *MYB33* and *MYB65* just to anthers (Alonso-Peral et al., 2010). Over-expressing *miR159* in Arabidopsis decreases the levels of MYB33 causing defects in anther development and male sterility (Achard et al., 2004; Schwab et al., 2005). In rice as in Arabidopsis, *OsGAMYB* expression is also restricted by *miR159* to anthers (Aya et al., 2009; Tsuji et al., 2006). *OsGAMYB* loss-of-function mutants have defective anthers and pollen (Kaneko et al., 2004).

The *miR319* family is encoded by the *MIR319a-c* loci and targets a subset of *TCP* transcription factor genes (*TCP2*, *TCP3*, *TCP4*, *TCP10* and *TCP24*) that are involved in multiple aspects of plant growth, including flower production, leaf and gametophyte development (Schommer et al., 2012). Over-expression of *miR319* affects leaf and cotyledon development, but also causes stamen defects and male sterility similar to what is observed in *miR159* over-expressing plants (Palatnik et al., 2007). A loss-of-function *miR319* mutant exhibited defects in petal and stamen development, such as narrower and shorter petals and abnormal anther formation (Nag et al., 2009).

The defects in the maturation of sepals, petals and anthers that are observed in plants where *miR159* and *miR319* activities are reduced through expression of target mimics resemble the defects observed in *auxin response factor 6 (arf6) arf8* double mutants (Rubio-Somoza and Weigel, 2011). *ARF6* and *ARF8* regulate the expression of auxin homeostatic genes and are involved in limiting the extent of cytokinin activity in the meristem and thus some of the floral defects in the meristems of *arf6/8* double mutants can be attributed to reduced auxin activity and increased cytokinin activity in the meristem (Rubio-Somoza and Weigel, 2011).

In Arabidopsis the patterns of expression of *ARF6* and *ARF8* in specific floral organs are regulated by *miR167* (Wu et al., 2006), and increased *miR167* levels results in the same floral phenotypes as reduced *ARF6/8* expression, ie. under-developed floral organs and reduced fertility (Ru et al., 2006). The expression of *miR167* is up-regulated independently by both TCP4 and MYB33 which are themselves regulated by *miR319* and *miR159* respectively (Fig. 3). Thus the overlapping functions of *miR319* and *miR159* can be attributed to the interaction between their own targets (TCP and MYB) as well as a common downstream target in *miR167* and its target genes *ARF6* and *ARF8* (Rubio-Somoza and Weigel, 2011). There is added complexity due to the fact that different isoforms of *miR67* differ in their expression patterns and in their ability to repress *ARF6/8* expression, and also due to the fact that there is cross-regulation between *miR167* genes such as between *miR167a* and *miR167c* (Rubio-Somoza and Weigel, 2011).

#### The role of *miR160* in flower development

*MiR160* is involved in the regulation of other *ARF* genes, it down-regulates the *ARF10*, *ARF16* and *ARF17* genes (Mallory et al., 2005; Liu et al., 2010). In the *floral organs in carpel (foc)* mutant there is a transposon insertion in the 3' regulatory region of *miR160* which results in reduced expression of this miRNA in flowers. As a result *ARF10*, *ARF16* and *ARF17* expression levels are elevated in the *foc* mutant and the mutant displays defects in floral organ formation such as reduced fertility, the appearance of floral organs inside siliques, and irregular flower shape, as well as

aberrant seeds, and viviparous seedlings (Liu et al., 2010). The *foc* mutant is deficient in its response to auxin as might be expected as ARF10, ARF16 and ARF17 are involved in auxin signaling. Interestingly, however, auxin was also found to control the levels of *miR160a* and up-regulates its expression through auxin response elements in the 3' regulatory region of the *MIR160* gene (Liu et al., 2010).

#### The role of *miR165* and *miR166* in flower development

In *Arabidopsis* *miR165* and *miR166* differ by only one nucleotide and both target the same *HD-ZIP III* genes *ATHB15*, *ATHB8*, *REVOLUTA*, *PHABULOSA*, and *PHAVOLUTA* (Floyd and Bowman, 2004; Reinhart et al., 2002; Zhou et al. 2007), although they exhibit distinct temporal and spatial expression patterns suggesting that they might regulate these target genes in different ways (Jung and Park 2007). Over-expression of *miR165* in the *Arabidopsis meristem enlargement 1 (men1)* mutant results in reduced the levels of all of these *HD-ZIP III* target genes and causes developmental defects in the SAM such as an enlarged apical meristem and short and sterile carpels (Kim et al., 2005). *MiR166* has also been shown to control embryonic SAM development in rice and maize (Nagasaki et al., 2007; Nogueira et al., 2007).

*MiR165* and *miR166* act in parallel to the *WUSCHEL-CLAVATA* pathway to regulate SAM development (Jung and Park 2007) and they do this through an interaction with ARGONAUTE 10 (AGO 10) (Zhu et al., 2011). Whilst *miR165/166* would normally bind to AGO 1 to form an active RNA-induced silencing complex (RISC), AGO 10 specifically interacts with *miR165/166* with a higher binding affinity than AGO 1. The

expression of AGO 10 in the SAM thus sequesters the *miR165/166* and prevents them forming an active RISC with AGO 1, thus *HD-ZIP III* expression levels are not repressed allowing normal SAM development (Zhu et al., 2011).

## **Sex determination in flowers**

Maize is monoecious, with male tassels and female ears. Flowers start as bisexual but later undergo stamen arrest in the ear, and pistil abortion in the tassel. In tassels of the recessive *tasselseed4* (*ts4*) and the dominant *Tasselseed6* (*Ts6*) mutants, pistils fail to abort and male floral organs do not develop, this results in pistils rather than stamens in the tassel. Cloning of the *Ts4* gene showed it to be an miRNA of the *miR172* gene family, called *zma-MIR172e*. Sequencing of *zma-MIR172e* from different *ts4* mutants demonstrated that insertion mutations in the promoter region of *zma-MIR172e* were responsible for the *ts4* mutant phenotypes. A target of *zma-miR172e* is the AP2-like gene *INDETERMINATE SPIKELET1* (*IDS1*), mutation of which partially suppress the *ts4* phenotype (Chuck 2007). Cloning of *Ts6* revealed it to be a mutated form of the *IDS1* gene, the mutation being in the *miR172* binding site thus preventing regulation by *zma-miR172e* which explains the similar phenotypes of the *ts4* and *Ts6* mutants. De-regulated expression of *IDS1* in these mutants represses the expression of MADS box genes in the maize spikelet, resulting in floral meristem indeterminacy and a failure of carpel abortion (Chuck 2007).

Interestingly, as Chuck et al. point out, the wheat domestication gene *Q* is an AP2-like gene orthologous to *IDS1*. The mutation that gave rise to the dominant *Q* allele that is

found in cultivated wheats occurred only once and is a C-to-T mutation at the 3' end of the *miR172* binding site (Chuck et al., 2007), which could explain its higher expression levels than the recessive *q* allele (Simons et al., 2006). It is possible that other mutations, similar to *ts4*, *Ts6* and *Q*, resulting in altered miRNA regulation of *AP2*-like genes involved in inflorescence development could also have had a role in the domestication of other crop species.

### **Biotechnological applications of miRNAs in manipulating the flowering processes**

MiRNAs offer the potential to the control of almost every aspect of plant development. Altering the expression levels of specific plant miRNAs, as well as the use of artificial miRNAs, makes it possible to regulate key transcription factors and thus entire downstream gene regulatory networks (Schwab et al., 2006). Economically important processes such as plant growth and stature, flowering, seed set and yield are obvious targets for the use of this technology to improve agricultural and horticultural productivity, there is also evidence that miRNAs regulate nodulation (Yan et al., 2013).

#### **Regulating flowering time**

Flowering time is an important target for plant breeding as it affects flower, seed and fruit development, ease of harvest and marketing (Jung and Muller 2009). Flowering in those crops and forages where the vegetative tissues are the important parts that are harvested has negative impacts on yield and nutritional quality, therefore inhibiting



flowering in these crops would be beneficial (although flowering is needed to enable seed production). On the other hand being able to induce early flowering would be of benefit to breeding companies enabling them to speed up their breeding programmes and develop new varieties more quickly, this is particularly true in tree species where it would be necessary to overcome their naturally long juvenile phases.

The activity of miRNAs can be reduced by the expression of specific target mimics that sequester the miRNA (Franco-Zorrilla et al., 2007). Conversely it is possible to increase miRNA activity generally throughout the plant, or in specific tissues, by overexpressing *MIR* genes or artificial miRNAs (amiRNAs) using constitutive or tissue-specific promoters (Li et al., 2013; Schwab et al., 2006). Schwab et al. expressed amiRNAs directed against different sequences of the Arabidopsis *FT* gene, *amiR-ft-1* or *amiR-ft-2*, to down-regulate this gene to try to delay flowering in transgenic plants. *FT* mRNA levels were reduced below RT-PCR detection levels in the *amiR-ft* expressing plants and flowering of these lines was delayed to a similar extent as in *ft* null mutants. This inhibition of flowering was also shown to work when an *amiR-FT* construct was expressed under the control of the *SUC2* promoter which is expressed specifically in phloem companion cells (Mathieu et al. 2007).

The *FT* gene acts partially redundantly with a closely related and partially redundant paralog called *TWIN SISTER OF FT (TSF)*, the *ft tsf* double mutant is even later flowering than the *ft* mutant. It is possible to use amiRNAs to down-regulate more than one gene target if the gene targets share enough homologous sequence for an amiRNA binding site (Schwab et al., 2006). Mathieu et al. did this in the case of *FT* and *TSF* by designing an

amiRNA that would target a common sequence present in both genes. The transgenic Arabidopsis plants expressing this *amiR-FT/TSF* construct flowered as late as the *ft tsf* double mutant (Mathieu et al 2007). Caution needs to be taken with this approach as Schwab et al. have shown that the degree of down-regulation of different target genes by an amiRNA can vary, and this variation is not correlated with the degree of complementarity to the target, or to the level of endogenous expression of the target gene (Schwab et al., 2006).

Li et al. showed that it was possible to both delay and advance the onset of flowering in ornamental Gloxinia (*Sinningia speciosa*) by up- or down-regulating the levels of *miR159*, which inhibits flowering by reducing the expression levels of the Gloxinia homologues of *GAMYB* and *LFY*. Transgenic lines with increased levels of *miR159* were delayed in the onset of flowering, whilst in those lines that had reduced levels of *miR159* caused by over-expression of a target mimic of *miR159* (*MIM159*), an early flowering phenotype was observed (Li et al., 2013). In a few transgenic lines where *MIM159* expression was very high, and where there was consequently strong repression of *miRNA159* and high levels of *LFY* expression, some petal to sepal conversion was observed. Thus it is not desirable to severely/completely inhibit the expression of *miR159*. Using inducible expression systems to regulate the levels of this miRNA could reduce this problem and also have the potential to control the timing of flowering in a very precise manner.

Such an inducible expression approach has been shown to work by Yeoh et al. (2011). They used Arabidopsis plants that were late flowering because they were expressing *amiR-FT* directed against the Arabidopsis *FT* gene (originally described in

Mathieu et al. 2007), and they transformed these plants with the *FTa1* gene from *Medicago truncatula*, an orthologous gene that has diverged sufficiently from the Arabidopsis equivalent to avoid suppression by the *amiR-FT*. The expression of the *FTa1* gene from Medicago in the *amiR-FT* Arabidopsis plants rescued the late flowering phenotype. Using an alcohol-inducible promoter to drive the expression of the *FTa1* gene enabled them to induce flowering in the transgenic plants in a fairly precise and synchronous manner by exogenous application of ethanol (Yeoh et al., 2011).

### Male sterility systems

In addition to manipulating the time of flowering, miRNAs are involved in male sterility (Achard et al. 2004; Toppino et al., 2011; Zhou et al., 2012) and can be used to create male sterile lines for F1 hybrid production. F1 hybrid production is mainly based on the three-line and two-line systems. The three-line system uses cytoplasmic male sterility lines, maintainer lines and restorer lines. The two-line hybrid system is based on environmentally sensitive (eg. photoperiod-sensitive (PGMS), or thermo-sensitive, (TGMS)) male sterile lines, which serve as both the male sterile lines and maintainer lines under different environmental conditions (Chen and Liu 2014). The two-line system is much simpler for breeding purposes and makes it easier to exploit hybrid vigour (heterosis).

PGMS and TGMS are widely used in two-line hybrid rice production but the molecular basis underlying this male sterility was unknown. Zhou et al. (2012) cloned the locus responsible for PGMS in a *japonica* rice line, and TGMS in an *indica* rice line (the same locus was responsible for both), and found that it encoded a novel small 21-nt RNA named osa-smR5864w, which was preferentially expressed in young panicles. The PGMS

and TGMS phenotypes in the male sterile lines was due to a C-to-G point mutation in the RNA, the mutated RNA being named osa-smR5864m. Expression of the wild-type miRNA, osa-smR5864w, restored fertility in both the PGMS and TGMS male sterile lines (Zhou et al. 2012). If this small RNA is conserved in other crop plants then it would be a clear target for site-directed mutagenesis to engineer PGMS or TGMS in those crop plants.

Another approach to engineer reversible male sterility using miRNAs was described by Toppino et al. (2011). In this approach anther-specific promoters were used to drive the expression of amiRNAs designed to target and repress the expression of factors involved in the basal RNA polymerase II transcription complex. Some of these factors are essential for cell viability and are highly conserved, Toppino et al. targeted two of these factors for repression by amiRNAs and by expressing the amiRNAs only in the tapetum or microspores they were able to generate lines in which pollen viability was greatly reduced. Reversibility of the male sterile phenotype was achieved through expression of amiRNA-insensitive forms of the target genes that had been designed with mis-matches in the amiRNA target sequence to make them insensitive to the activity of the amiRNAs. Expressing these amiRNA-insensitive alleles from an alcohol-inducible promoter enabled fertility in these male-sterile lines to be restored by ethanol treatment.

In addition to regulating flowering time and the development of male sterility systems in different crops as outlined above, which have probably the most obvious economic benefits, other potential flowering-related targets for manipulation by miRNA approaches could be altering flower shape/appearance, and possibly even controlling the sex of the flowers in monoecious and dioecious plant species (Chuck et al., 2007; Song et

al. 2013). As new discoveries reveal even more roles for miRNAs in flowering-related processes, perhaps some specific to individual species, then the potential biotechnological applications will only increase further.

## Bibliography

- Achard, P., Herr, A., Baulcombe, D. C., & Harberd, N. P. (2004) Modulation of floral development by a gibberellin-regulated microRNA. *Development* **131**(14), 3357-3365.
- Adam, H., Marguerettaz, M., Qadri, R., Adroher, B., Richaud, F., Collin, M., Thuillet, A., Vigouroux, Y., Laufs, P., Tregear, J.W. and Jouannic, S. (2011) Divergent expression patterns of miR164 and CUP-SHAPED COTYLEDON genes in palms and other monocots: implication for the evolution of meristem function in angiosperms. *Mol. Biol. Evol.* **28**, 1439–1454.
- Allen, R.S., Li, J.Y., Stahle, M.I., Dubroue, A., Gubler, F. and Millar, A.A. (2007) Genetic analysis reveals functional redundancy and the major targets of the *Arabidopsis* miR159 family. *PNAS* **104**, 16371-16376.
- Alonso-Peral, M.M., Li, J., Li, Y., Allen, R.S., Schnippenkoetter, W., Ohms, S., White, R.G. and Millar, A.A. (2010) The microRNA159-regulated GAMYB-like genes inhibit growth and promote programmed cell death in *Arabidopsis*. *Plant Physiology* **154**, 757-771.
- Amasino, R. (2010) Seasonal and developmental timing of flowering. *Plant J.* **61**, 1001–1013
- Aukerman, M.J., and Sakai, H. (2003) Regulation of flowering time and floral organ identity by a microRNA and its APETALA2-like target genes. *Plant Cell* **15**, 2730-2741.
- Aya, K., Ueguchi-Tanaka, M., Kondo, M., Hamada, K., Yano, K., Nishimura, M. and Matsuoka, M., (2009) Gibberellin modulates anther development in rice via the

- transcriptional regulation of GAMYB. *Plant Cell* **21**,1453–1472.
- Baker, C.C., Sieber, P., Wellmer, F. and Meyerowitz, E.M. (2005) The early extra petals1 mutant uncovers a role for microRNA miR164c in regulating petal number in *Arabidopsis*. *Curr. Biol.*, **15**, 303–315.
- Berger, Y., Harpaz-Saad, S., Brabdt, A., Melnik, H., Sirding, N., Alvarez, J.P., Zinder, M., Samach, A., Eshed, Y. and Ori, N. (2009) The NAC-domain transcription factor GOBLET specifies leaflet boundaries in compound tomato leaves. *Development*, **136**, 823–832.
- Bergonzi, S., Albani, M.C., Ver Loren van Themaat E., Nordström, K.J., Wang, R., Schneeberger, K., Moerland, P.D. and Coupland, G. (2013) Mechanisms of age-dependent response to winter temperature in perennial flowering of *Arabis alpina*. *Science*, **340**, 1094-1097.
- Cartolano, M., Castillo, R., Efremova, N., Kuckenberg, M., Zethof, J., Gerats, T., Schwarz-Sommer, Z., Vandenbussche, M. (2007) A conserved microRNA module exerts homeotic control over *Petunia hybrida* and *Antirrhinum majus* floral organ identity. *Nat. Genet.* **39**, 901–905.
- Causier, B., Schwarz-Sommer, Z. and Davies, B. (2010) Floral organ identity: 20 years of ABCs. *Semin. Cell Dev. Biol.*, **21**, 73–79.
- Chen X. (2004) A microRNA as a translational repressor of APETALA2 in *Arabidopsis* flower development. *Science*, **303**, 2022-2025.
- Chen, L., & Liu, Y. G. (2014) Male sterility and fertility restoration in crops. *Annual review of plant biology*, **65**, 579-606.

- Chuck, G., Meeley, R., Irish, E., Sakai, H., & Hake, S. (2007) The maize tasselseed4 microRNA controls sex determination and meristem cell fate by targeting Tasselseed6/indeterminate spikelet1. *Nature genetics*, **39**, 1517-1521.
- Chuck, G., Candela, H., & Hake, S. (2009) Big impacts by small RNAs in plant development. *Current opinion in plant biology*, **12**, 81-86.
- Chiou, T.J., Aung, K., Lin, S.I., Wu, C.C., Chiang, F., and Su, C.L. (2006) Regulation of phosphate homeostasis by microRNA in *Arabidopsis*. *Plant Cell*, **18**, 412-421.
- Endo, Y., Iwakawa, H.O. and Tomari, Y. (2013) *Arabidopsis* ARGONAUTE7 selects miR390 through multiple checkpoints during RISC assembly. *EMBO Reports* **14**, 652-658.
- Fahlgren, N., Montgomery, T.A., Howell, M.D., Allen, E., Dvorak, S.K., Alexander, A.L. and Carrington, J.C. (2006) Regulation of AUXIN RESPONSE FACTOR3 by TAS3 ta-siRNA affects developmental timing and patterning in *Arabidopsis*. *Current Biology* **16**, 939-944.
- Floyd, S. K., & Bowman, J. L. (2004) Gene regulation: ancient microRNA target sequences in plants. *Nature*, **428**, 485-486.
- Fornara, F., de Montaigu, A. and Coupland, G. (2010) SnapShot: control of flowering in *Arabidopsis*. *Cell* **141**, 550-550.e2.
- Franco-Zorrilla, J.M., Valli, A., Todesco, M., Mateos, I., Rubio-Somoza, I., Leyva, A., Weigel, D., Garcia, J.A. and Paz-Ares, J. (2007) Target mimicry provides a new mechanism for regulation of microRNA activity. *Nature Genetics* **39**, 1033-1037.
- Galvão, V. C., Horrer, D., Küttner, F., & Schmid, M. (2012) Spatial control of flowering by DELLA proteins in *Arabidopsis thaliana*. *Development*, 4072-4082.



- Garcia D. (2008) A miracle in plant development: Role of MicroRNAs in cell differentiation and patterning. *Seminars in Cell & Developmental Biology* **19**, 586-595.
- Grigorova, B., Mara1, C., Hollender, C., Sijacic, P., Chen, X. and Liu, Z. (2011) LEUNIG and SEUSS co-repressors regulate miR172 expression in Arabidopsis flowers. *Development* **138**, 2451–2456.
- Guo, A.Y., Zhu, Q.H., Gu, X., Ge, S., Yang, J. and Luo J. (2008) Genome-wide identification and evolutionary analysis of the plant specific SBP-box transcription factor family. *Gene* **418**, 1-8.
- Huijser, P. and Schmid, P. (2011) The control of developmental phase transitions in plants. *Development* **138**, 4117-4129.
- Jackson, S.D. (2009) Plant responses to photoperiod. *New Phytologist* **181**, 517-531.
- Jin, D., Wang, Y., Zhao, Y. and Chen, M. (2013) MicroRNAs and their cross-talks in plant development. *Journal of Genetics and Genomics* **40**, 161-170.
- Jones-Rhoades, M.W. and Bartel, D.P. (2004) Computational identification of plant microRNAs and their targets, including a stress-induced miRNA. *Molecular Cell* **14**, 787–799.
- Jones-Rhoades, M., Bartel, D.P. and Bartel, B. (2006) MicroRNAs and their regulatory roles in plants. *Annual Review of Plant Biology* **57**, 19-53.
- Jung, J.H., Seo, Y.H., Seo, P.J., Reyes, J.L., Yun, J., Chua, N.H. and Park, C.M. (2007) *Plant Cell* **19**, 2736-48.
- Jung, C. and Muller, A.E. (2009) Flowering time control and applications in plant breeding. *Trends in Plant Science*, **14**, 563-573.

- Jung, J. H., & Park, C. M. (2007) MIR166/165 genes exhibit dynamic expression patterns in regulating shoot apical meristem and floral development in Arabidopsis. *Planta*, **225**, 1327-1338.
- Kaneko, M., Inukai, Y., Ueguchi-Tanaka, M., Itoh, H., Izawa, T., Kobayashi, Y., Hattori, T., Miyao, A., Hirochika, H., Ashikari, M. and Matsuoka, M. (2004) Loss-of-function mutations of the rice GAMYB gene impair  $\alpha$ -amylase expression in aleurone and flower development. *Plant Cell* **16**, 33–44.
- Kasai, A., Kanehira, A., & Harada, T. (2010) miR172 can move long distances in *Nicotiana benthamiana*. *Open Plant Sci. J*, **4**, 1-6.
- Kim, J., Jung, J. H., Reyes, J. L., Kim, Y. S., Kim, S. Y., Chung, K. S., Kim, J., Lee, M., Lee, Y., Kim, V.N., Chua, N-H. & Park, C. M. (2005) microRNA- directed cleavage of ATHB15 mRNA regulates vascular development in Arabidopsis inflorescence stems. *The Plant Journal*, **42**, 84-94.
- Kim, W., Ahn, H.J., Chiou, T.J. and Ahn, J.H. (2011) The role of the miR399-PHO2 module in the regulation of flowering time in response to different ambient temperatures in *Arabidopsis thaliana*. *Molecules and Cells*, **32**, 83-88.
- Kim, J.J, Lee, J.H., Kim, W., Jung, H.S., Huijser, P. and Ahn, J.H. (2012) The microRNA156-SQUAMOSA PROMOTER BINDING PROTEIN-LIKE3 module regulates ambient temperature-responsive flowering via FLOWERING LOCUS T in *Arabidopsis*. *Plant Physiology* **159**, 461-478.
- Kruszka, K., Pieczynski, M., Windels, D., Bielewicz, D., Jarmolowski, A., Szweykowska-Kulinska, Z. and Vazquez, F. (2012) Role of microRNAs and other sRNAs of plants in their changing environments. *Journal of Plant Physiology* **169**, 1664-1672.

- Laufs, P., Peaucelle, A., Morin, H. and Traas, J. (2004) MicroRNA regulation of the CUC genes is required for boundary size control in *Arabidopsis* meristems. *Development*, **131**, 4311–4322.
- Lee, H., Yoo, S.J., Lee, J.H., Kim, W., Yoo, S.K., Fitzgerald, H., Carrington, J.C. and Ahn, J.H. (2010) Genetic framework for flowering-time regulation by ambient temperature-responsive miRNAs in *Arabidopsis*. *Nucleic Acids Research*, **38**, 3081- 3093.
- Lee, D.Y. and An, G. (2012) Two AP2 family genes, SUPERNUMERARY BRACT (SNB) and OsINDETERMINATE SPIKELET1 (OsIDS1), synergistically control inflorescence architecture and floral meristem establishment in rice. *Plant J.* **69**, 445–461.
- Li, Y., Fu, Y., Ji, L., Wu, C., & Zheng, C. (2010) Characterization and expression analysis of the *Arabidopsis* mir169 family. *Plant science*, **178**, 271-280.
- Li, Y., Li, C., Ding, G. and Jin, Y. (2011) Evolution of MIR159/319 microRNA genes and their post-transcriptional regulatory link to siRNA pathways. *BMC Evol. Biol.* **11**, 122.
- Li, X., Bian, H., Song, D., Ma, S., Han, N., Wang, J. and Zhu, M. (2013) Flowering time control in ornamental gloxinia (*Sinningia speciosa*) by manipulation of miR159 expression. *Annals of Botany*. **111**, 791-799.
- Liu, X., Huang, J., Wang, Y., Khanna, K., Xie, Z., Owen, H. A., & Zhao, D. (2010) The role of floral organs in carpels, an *Arabidopsis* loss- of- function mutation in MicroRNA160a, in organogenesis and the mechanism regulating its expression. *The Plant Journal*, **62**, 416-428.
- Liu, J. and Vance, C.P. (2010) Crucial roles of sucrose and microRNA399 in systemic signaling of P deficiency: a tale of two team players? *Plant Signaling & Behaviour*, **5**, 1556-1560.

- Luo, Y., Guo, Z., & Li, L. (2013) Evolutionary conservation of microRNA regulatory programs in plant flower development. *Developmental biology*, **380**, 133-144.
- Mallory, A.C., Bartel,D.P. and Bartel,B., (2005) MicroRNA-directed regulation of Arabidopsis AUXIN RESPONSE FACTOR 17 is essential for proper development and modulates expression of early auxin response genes. *Plant Cell* **17**, 1360–1375.
- Martin, A., Adam, H., Díaz-Mendoza, M., Żurczak, M., González-Schain, N. D. & Suárez-López, P. (2009) Graft-transmissible induction of potato tuberization by the microRNA *miR172*. *Development*, **136**, 2873-81.
- Mathieu, J., Warthmann, N., Küttner, F. & Schmid, M. (2007) Export of FT Protein from Phloem Companion Cells Is Sufficient for Floral Induction in Arabidopsis. *Current Biology*, **17**, 1055-1060.
- Mathieu, J., Yant, L.J., Mürdter, F., Küttner, F. and Schmid. M. (2009) Repression of flowering by the miR172 target SMZ. *PLoS Biology* **7**, e1000148.
- Matsoukas, I.G., Massiah, A.J. and Thomas, B. (2012) Florigenic and antiflorigenic signaling in plants. *Plant Cell Physiology* **53**, 1827-1842.
- Montgomery, T.A., Howell, M.D., Cuperus, J.T., Li, D., Hansen, J.E., Alexander, A.L., Chapman, E.J., Fahlgren, N., Allen, E. and Carrington, J.C. (2008) Specificity of ARGONAUTE7-miR390 interaction and dual functionality in TAS3 trans-acting siRNA formation. *Cell* **133**, 128-141.
- Nair, S.K., Wang, N., Turuspekov, Y., Pourkheirandish, M., Sinsuwongwat, S., Chen, G., Sameri, M., Tagiri, A., Honda, I., Watanabe, Y., Kanamori, H., Wicker, T., Stein, N., Nagamura, Y., Matsumoto, T. and Komatsuda, T. (2010) Cleistogamous flowering in

barley arises from the suppression of microRNA-guided HvAP2 mRNA cleavage.

*PNAS*, **107**, 490–495.

Nag, A., King, S. and Jack, T. (2009) miR319a targeting of TCP4 is critical for petal growth and development in Arabidopsis. *PNAS*, **106**, 22534–22539.

Nagasaki, H., Itoh, J. I., Hayashi, K., Hibara, K. I., Satoh-Nagasawa, N., Nosaka, M., Mukouhata, M., Ashikari, M., Kitano, H., Matsuoka, M., Nagato, Y., & Sato, Y. (2007) The small interfering RNA production pathway is required for shoot meristem initiation in rice. *PNAS*, **104**, 14867-14871.

Nogueira, F. T., Madi, S., Chitwood, D. H., Juarez, M. T., & Timmermans, M. C. (2007) Two small regulatory RNAs establish opposing fates of a developmental axis. *Genes & development*, **21**, 750-755.

Palatnik, J. F., Wollmann, H., Schommer, C., Schwab, R., Boisbouvier, J., Rodriguez, R., Warthmann, N., Allen, E., Dezulian, T., Huson, D., Carrington, J.J. and Weigel, D. (2007) Sequence and expression differences underlie functional specialization of *Arabidopsis* microRNAs miR159 and miR319. *Developmental Cell*, **13**, 115-125.

Poethig, R. S. (2003) Phase change and the regulation of developmental timing in plants. *Science*, **301**, 334–336

Posé, D., Yant, L. and Schmid, M. (2012) The end of innocence: flowering networks explode in complexity. *Curr. Opin. Plant Biol.*, **15**, 45–50

Reinhart, B. J., Weinstein, E. G., Rhoades, M. W., Bartel, B., & Bartel, D. P. (2002) MicroRNAs in plants. *Genes & development*, **16**, 1616-1626.

Ru, P., Xu, L., Ma, H., & Huang, H. (2006) Plant fertility defects induced by the enhanced expression of microRNA167. *Cell research*, **16**, 457-465.

- Rubio-Somoza, I. and Weigel, D. (2011) MicroRNA networks and developmental plasticity in plants. *Trends in Plant Science*, **16**, 258-264.
- Rubio-Somoza, I. and Weigel, D. (2013) Coordination of flower maturation by a regulatory circuit of three microRNAs. *PLoS genetics*, **9**, e1003374.
- Schommer, C., Bresso, E.G., Spinelli, S.V. and Palatnik, J.F. (2012) Role of microRNA miR319 in plant development. *Signaling and Communication in Plants*, **15**, 29-47.
- Schwab, R., Palatnik, J.F., Riester, M., Schommer, C., Schmid, M. and Weigel, D. (2005) Specific effects of microRNAs on the plant transcriptome. *Dev. Cell*, **8**, 517–527.
- Schwab, R., Ossowski, S., Riester, M., Warthmann, N. and Weigel, D. (2006) Highly specific gene silencing by artificial microRNAs in *Arabidopsis*. *Plant Cell*, **18**, 1121-1133.
- Schwarz, S., Grande, A.V., Bujdosó, N., Saedler, H., and Huijser, P. (2008) The microRNA regulated SBP-box genes SPL9 and SPL15 control shoot maturation in *Arabidopsis*. *Plant Molecular Biology* **67**, 183-195.
- Sgamma, T., Jackson, A., Muleo, R., Thomas, B., & Massiah, A. (2014) TEMPRANILLO is a regulator of juvenility in plants. *Scientific reports*, **4**.
- Simons, K. J., Fellers, J. P., Trick, H. N., Zhang, Z., Tai, Y. S., Gill, B. S., & Faris, J. D. (2006) Molecular characterization of the major wheat domestication gene Q. *Genetics*, **172**, 547-555.
- Sieber, P., Wellmer, F., Gheyselinck, J., Riechmann, J.L. and Meyerowitz, E.M. (2007) Redundancy and specialization among plant microRNAs: role of the MiR164 family in developmental robustness. *Development*, **134**, 1051–1060.
- Song, Y., Ma, K., Ci, D., Zhang, Z. and Zhang, D. (2013) Sexual Dimorphism Floral

- MicroRNA Profiling and Target Gene Expression in Andromonoecious Poplar (*Populus tomentosa*) *PLoS ONE*, **8**, e62681.
- Spanudakis, E. and Jackson, S. (2014) The role of microRNAs in the control of flowering time. *Journal of Experimental Botany*, **65**, 365-380.
- Srikanth, A. and Schmid, M. (2011) Regulation of flowering time: all roads lead to Rome. *Cellular and Molecular Life Sciences*, **68**, 2013-2037.
- Sunkar, R. and Jagadeeswaran, G. (2008) In silico identification of conserved microRNAs in large number of diverse plant species. *BMC Plant Biol.*, **8**, 37.
- Takada, S., Hibara, K., Ishida, T. and Tasaka. (2001) The CUP-SHAPED COTYLEDON1 gene of *Arabidopsis* regulates shoot apical meristem formation. *Development*, **128**, 1127–1135.
- Terzi, L.C. and Simpson, G.G. (2008) Regulation of flowering time by RNA processing. *Current Topics in Microbiology and Immunology*, **326**, 201-218.
- Todesco, M., Rubio-Somoza, I., Paz-Ares, J., & Weigel, D. (2010) A collection of target mimics for comprehensive analysis of microRNA function in *Arabidopsis thaliana*. *PLoS genetics*, **6**, e1001031.
- Toppino, L., Kooiker, M., Lindner, M., Dreni, L., Rotino, G. L., & Kater, M. M. (2011) Reversible male sterility in eggplant (*Solanum melongena* L.) by artificial microRNA-mediated silencing of general transcription factor genes. *Plant biotechnology journal*, **9**, 684-692.
- Tsuji, H., Aya, K., Ueguchi-Tanaka, M., Shimada, Y., Nakazono, M., Watanabe, R., Nishizawa, N. K., Gomi, K., Shimada, A., Kitano, H., Ashikari, M. and Matsuoka, M.

- (2006) GAMYB controls different sets of genes and is differentially regulated by microRNA in aleurone cells and anthers. *Plant J.*, **47**, 427–444.
- Vroemen, C.W., Mordhorst, A.P., Albrecht, C., Kwaaitaal, M.A.C.J. and de Vries, S.C. (2003) The CUP-SHAPED COTYLEDON3 gene is required for boundary and shoot meristem formation in *Arabidopsis*. *Plant Cell*, **15**, 1563–1577.
- Wahl, V., Ponnu, J., Schlereth, A., Arrivault, S., Langenecker, T., Franke, A., Feil, R., Lunn, J.E., Stitt, M. and Schmid, M. (2013) Regulation of flowering by trehalose-6-phosphate signaling in *Arabidopsis thaliana*. *Science*, **339**, 704–707.
- Wang, J.W. (2014) Regulation of flowering time by the miR156-mediated age pathway. *J. Exp. Bot.*, **65**, 4723–4730
- Wang, J.W., Czech, B. and Weigel, D. (2009) MiR156-regulated SPL transcription factors define an endogenous flowering pathway in *Arabidopsis thaliana*. *Cell*, **138**, 738–749.
- Willmann, M.R. and Poethig, R.S. (2007) Conservation and evolution of miRNA regulatory programs in plant development. *Curr. Opin. Plant Biol.* **10**, 503–511.
- Wollmann, H., Mica, E., Todesco, M., Long, J.A. and Weigel, D. (2010) On reconciling the interactions between APETALA2, miR172 and AGAMOUS with the ABC model of flower development. *Development*, **137**, 3633–3642.
- Wu, G., & Poethig, R. S. (2006) Temporal regulation of shoot development in *Arabidopsis thaliana* by miR156 and its target SPL3. *Development*, **133**, 3539–3547.
- Wu, M. F., Tian, Q., & Reed, J. W. (2006) *Arabidopsis* microRNA167 controls patterns of ARF6 and ARF8 expression, and regulates both female and male reproduction. *Development*, **133**, 4211–4218.
- Wu, G., Park, Y., Conway, S.R., Wang, J.W., Weigel, D. and Poethig, R.S. (2009) The



- sequential action of miR156 and miR172 regulates developmental timing in *Arabidopsis*. *Cell*, **138**, 750-759.
- Xie, K., Wu, C. and Xiong, L. (2006) Genomic organization, differential expression, and interaction of SQUAMOSA promoter-binding-like transcription factors and microRNA156 in rice. *Plant Physiology*, **142**, 280-293.
- Xu, M. Y., Zhang, L., Li, W. W., Hu, X. L., Wang, M. B., Fan, Y. L., Zhang, C.Y. & Wang, L. (2013) Stress-induced early flowering is mediated by miR169 in *Arabidopsis thaliana*. *J. Exp. Bot.*, **65**, 89-101.
- Yeoh, C.C., Balcerowicz, M., Laurie, R., Macknight, R. and Putterill, J. (2011) Developing a method for customized induction of flowering. *BMC Biotechnology*, **11**, 1-11.
- Yamaguchi, A., Wu, M., Yang, L, Wu, G., Poethig, R.S. and Wagner, D. (2009) The microRNA-regulated SBP-box transcription factor SPL3 is a direct upstream activator of LEAFY, FRUITFULL and APETALA1. *Developmental Cell*, **17**, 268-278.
- Yamaguchi, A. and Abe, M. (2012) Regulation of reproductive development by non-coding RNA in *Arabidopsis*: to flower or not to flower. *Journal of Plant Research*, **125**, 693-704.
- Yan, Z., Hossain, M.S., Wang, J., Valdés-López, O., Liang, Y., Libault, M., Qiu, L. and Stacey, G. (2013) miR172 regulates soybean nodulation. *Molecular Plant-Microbe Interactions*, **26**, 1371-1377.
- Yang, L., Xu, M., Koo, Y., He, J. and Poethig, R.S. (2013) Sugar promotes vegetative phase change in *Arabidopsis thaliana* by repressing the expression of MIR156A and MIR156C. *eLife*, **2**.
- Yant, L., Mathieu, J., Dinh, T.T, Ott, F., Lanz, C, Wollmann, H., Chen, X. and Schmid, M.

- (2010) Orchestration of the floral transition and floral development in *Arabidopsis* by the bifunctional transcription factor APETALA2. *Plant Cell*, **22**, 2156-2170.
- Yu, S., Cao, L., Zhou, C.M., Zhang, T.Q., Lian, H., Sun, Y., Wu, J., Huang, J., Wang, G. and Wang, J.W. (2013) Sugar is an endogenous cue for juvenile-to-adult phase transition in plants. *eLife*, **2**.
- Yu, S., Galvão, V. C., Zhang, Y. C., Horrer, D., Zhang, T. Q., Hao, Y. H., Feng, Y.Q., Wang, S., Schmid, M. and Wang, J.W. (2012) Gibberellin regulates the Arabidopsis floral transition through miR156-targeted SQUAMOSA PROMOTER BINDING-LIKE transcription factors. *The Plant Cell Online*, **24**, 3320-3332.
- Zhang, X., Zou, Z., Zhang, J., Zhang, Y., Han, Q., Hu, T., Xu, X., Liu, H., Li, H. and Ye, Z. (2011) Over-expression of *sly-miR156a* in tomato results in multiple vegetative and reproductive trait alterations and partial phenocopy of the *sft* mutant. *FEBS Letters*, **21**, 435-439.
- Zhao, L., Kim, Y., Dinh, T.T. and Chen, X. (2007) MiR172 regulates stem cell fate and defines the inner boundary of APETALA3 and PISTILLATA expression domain in *Arabidopsis* floral meristems. *Plant J.*, **51**, 840–849.
- Zhao, B., Ge, L., Liang, R., Li, W., Ruan, K., Lin, H., & Jin, Y. (2009) Members of miR-169 family are induced by high salinity and transiently inhibit the NF-YA transcription factor. *BMC Molecular Biology*, **10**, 29.
- Zhou, G. K., Kubo, M., Zhong, R., Demura, T., & Ye, Z. H. (2007) Overexpression of miR165 affects apical meristem formation, organ polarity establishment and vascular development in *Arabidopsis*. *Plant and Cell Physiology*, **48**, 391-404.
- Zhou, C.M. and Wang, J.W. (2013) Regulation of flowering time by microRNAs. *Journal*

*of Genetics and Genomics*, **40**, 211-215.

Zhou, H., Liu, Q., Li, J., Jiang, D., Zhou, L., Wu, P., Sen, L., Feng, L., Zhu, L., Chen, L.,

Liu, Y-G. & Zhuang, C. (2012) Photoperiod-and thermo-sensitive genic male sterility in rice are caused by a point mutation in a novel noncoding RNA that produces a small RNA. *Cell research*, **22**, 649-660.

Zhu, Q.H., Upadhyaya, N.M., Gubler, F., Helliwell, C.A. (2009) Over-expression of miR172 causes loss of spikelet determinacy and floral organ abnormalities in rice (*Oryza sativa*). *BMC. Plant Biol.*, **9**, 17.

Zhu, Q.H. and Helliwell. C.A. (2011) Regulation of flowering time and floral patterning by miR172. *J. Exp. Bot.*, **62**, 487–495

Zhu, H., Hu, F., Wang, R., Zhou, X., Sze, S. H., Liou, L. W., Barefoot, A., Dickman, M, & Zhang, X. (2011) Arabidopsis Argonaute10 Specifically Sequesters miR166/165 to Regulate Shoot Apical Meristem Development. *Cell*, **145**, 242-256.

## Figures




Fig.1

The involvement of various miRNAs in the different flowering-related phases of plant development, from the transition from juvenility to maturity, to floral induction and development.




Fig.2

The changes associated with the transition from juvenility to maturity. MiR156 levels reduce with age influenced by sugar signaling pathways, this results in an increase in the levels of SPL proteins and miR172 which reduces the level of AP2-like floral repressors. The loss of the AP2-like repressors allows flowering to occur in inductive environmental conditions. AP2-like: APETELA 2-like, CO: CONSTANS, FLC: FLOWERING LOCUS C, SPL: SQUAMOSA PROMOTER BINDING-LIKE, T-6-P: TRHALOSE-6-PHOSPHATE.




Fig.3

An overview summarizing the action of miRNAs and their target genes in setting boundaries and regulating floral organ gene expression. MiR159 and miR319 and their target *GAMYB* and *TCP* proteins interact to regulate miR167 and the levels of ARF6 and ARF8. GA relieves the repression of both miR159 and *GAMYB* by DELLA proteins. *GAMYB* activates *LFY* to promote flowering, and possibly also might up-regulate miR159. AP2: APETELA 2, ARF: AUXIN RESPONSE FACTOR, CUC: CUP-SHAPED COTYLEDON, GA: Gibberellic acid, GOB: GOBLET, LFY: LEAFY, LUG: LEUNIG, PHAB: PHABULOSA, PHAV: PHAVOLUTA, NAM: NO APICAL MERISTEM, REV: REVOLUTA, SOC1: SUPPRESSOR OF CONSTANS 1